arXiv:gr-qc/0206029v2 24 Jun 2002

Dynamical N-body Equlibrium in Circular Dilaton Gravity

R. Kerner$^{(a,b)}$ and R. B. Mann$^{(a)}$
$^{(a)}$*Dept. of Physics, University of Waterloo, Waterloo, Ontario*
N2L 3G1, Canada
$^{(b)}$*Dept. of Physics, University of Alberta, Edmonton, Alberta*
V6T 1W5, Canada

We obtain a new exact equilibrium solution to the N-body problem in a one-dimensional relativistic self-gravitating system. It corresponds to an expanding/contracting spacetime of a circle with N bodies at equal proper separations from one another around the circle. Our methods are straightforwardly generalizable to other dilatonic theories of gravity, and provide a new class of solutions to further the study of (relativistic) one-dimensional self-gravitating systems.

Circular gravity is a general relativistic gravitational system in $(1+1)$ dimensions in which the topology of space is equivalent to that of a circle S^1. Its counterpart, lineal gravity (the same system with the S^1 topology replaced by $\mathbb{R}^1$), has been the subject of much investigation over the past twenty years, motivated by the desire to understand black holes [1], spacetime structure [2,3], canonical quantum gravity [4], string physics [5], information loss [6] and the N-body problem [7,8]. This latter problem involves consideration of a collection of N particles interacting through their own mutual gravitational attraction, along with other specified forces. Apart from functioning as prototypes for higher dimensional gravity, in lineal gravity such self-gravitating systems also approximate the behaviour of some physical systems in 3 spatial dimensions, such as stellar dynamics orthogonal to a galactic plane, the collisions of flat parallel domain walls, and the dynamics of cosmic strings. The ergodic and equipartition properties of non-relativistic one-dimensional self-gravitating systems are a current subject of research [9], and recently such systems were shown to exhibit fractal behaviour [10]. In the past few years these systems have been extended to relativistic lineal gravity. The 2-body problem has been solved exactly [11], and has been extended to include both cosmological expansion [12,13] and electromagnetic interactions [14]. The N-body extension of these systems, while considerably more complicated, is nevertheless amenable to analysis [8] and includes the non-relativistic self-gravitating systems [9,15] as a limiting case.

Comparitively little is known about the circular counterparts of these systems; they introduce qualitatively new features that are absent in the non-compact lineal case. For example, in the absence of other fields there are no static non-relativistic equilibrium solutions for N particles on a circle. This is because the non-relativistic potential grows linearly with increasing distance from the source, and a circular topology admits no solutions to the matching conditions for any $N \geq 1$.

We present here an exact equilibrium solution to the N-body problem in circular gravity. Our solution corresponds to N equal mass particles at equidistant proper separations from one another in a spacetime that expands/contracts in response to these sources. The configuration we find therefore corresponds to a dynamic, albeit unstable, equilibrium. To our knowledge these are the first N-body dynamic equilibrium solutions in any relativistic theory of gravity. While our solutions depend upon the choice of dilaton potential, our approach is straightforwardly generalizable to any dilatonic theory of gravity, and provides a new class of solutions to further the study of (relativistic) one-dimensional self-gravitating systems.

The action integral for a generic dilaton theory of gravity [4,16] minimally coupled to N point masses is

$$I = \int d^2x \left[\frac{1}{2\kappa}\sqrt{-g}g^{\mu\nu}\left\{ \Psi R_{\mu\nu} + \frac{1}{2}\nabla_\mu\Psi\nabla_\nu\Psi + g_{\mu\nu}V\left(\Psi\right)\right\} \right.$$
$$\left. + \sum_a \int d\tau_a \left\{ -m_a \left(-g_{\mu\nu}(x)\frac{dz_a^\mu}{d\tau_a}\frac{dz_a^\nu}{d\tau_a}\right)^{1/2} \right\} \delta^2(x - z_a(\tau_a)) \right] \ , \tag{1}$$

where Ψ is the dilaton field, $g_{\mu\nu}$ is the metric with determinant g and covariant derivative ∇_μ , R is the Ricci scalar, and τ_a is the proper time of a-th particle, with $\kappa = 8\pi G/c^4$. For circular gravity the range of the spatial coordinate x is $-L \leq x \leq L$, and the circular topology implies that all fields must be smooth (or at least C^1) functions of x with period $2L$, which implies

$$f(L) = f(-L) \qquad \text{and} \qquad f'(L) = f'(-L) = 0 \ . \tag{2}$$

for all functions. Henceforth we set $V\left(\Psi\right) = \Lambda/2$ throughout, where Λ is constant.

The action (1) describes a generally covariant self-gravitating system (without collisional terms, so that the bodies pass through each other); its field equations are

$$R - \Lambda = \kappa T^{P\mu}_{\ \ \mu} \qquad \frac{d}{d\tau_a}\left\{\frac{dz_a^\nu}{d\tau_a}\right\} + \Gamma^\nu_{\alpha\beta}(z_a)\frac{dz_a^\alpha}{d\tau_a}\frac{dz_a^\beta}{d\tau_a} = 0 \tag{3}$$

$$\frac{1}{2}\nabla_\mu\Psi\nabla_\nu\Psi - g_{\mu\nu}\left(\frac{1}{4}\nabla^\lambda\Psi\nabla_\lambda\Psi - \nabla^2\Psi\right) - \nabla_\mu\nabla_\nu\Psi = \kappa T^P_{\mu\nu} + \frac{\Lambda}{2}g_{\mu\nu} \tag{4}$$

where the stress-energy due to the point masses is

$$T^P_{\mu\nu} = \sum_{a=1}^{N} m_a \int d\tau_a \frac{1}{\sqrt{-g}} g_{\mu\sigma}g_{\nu\rho}\frac{dz_a^\sigma}{d\tau_a}\frac{dz_a^\rho}{d\tau_a}\delta^2(x - z_a(\tau_a)) \tag{5}$$

and is conserved. The preceding system generalizes Jackiw-Teitelboim lineal gravity [18], which equates scalar curvature to a (cosmological) constant Λ, reducing to it if the stress-energy vanishes. The specific choice of dilaton potential ensures that the evolution of the dilaton does not modify the aformentioned reciprocal gravity/matter dynamics. The system (3) is a closed system of $N+1$ that describes the evolution of the single metric degree of freedom and the N degrees of freedom of the point masses. The evolution of the point-masses governs the evolution of the dilaton via (4), whose divergencelessness is consistent with the conservation of $T^P_{\mu\nu}$, yielding only one independent equation to determine the single degree of freedom of the dilaton.

Making use of the previously formulated general framework for canonical circular dilaton gravity [17], we write the metric as $ds^2 = -\left(N_0 dt\right)^2 + \gamma\left(dx + \frac{N_1}{\gamma}dt\right)^2$. The extrinsic curvature K is then

$$K = (2N_0\gamma)^{-1}(2\partial_1 N_1 - \gamma^{-1}N_1\partial_1\gamma - \partial_0\gamma) = \sqrt{\gamma}\kappa(\pi - \Pi/\gamma) \tag{6}$$

where the conjugate momenta to γ and Ψ are respectively denoted π and Π. We can take the extrinsic curvature K to be a time coordinate $\tau(t)$ of the circle, thereby allowing the elimination of π from all field equations and the identification of the Hamiltonian from the action (1) after canonical reduction [17]:

$$I = \int d^2x\left\{\sum_a p_a\dot{z}_a\delta(x - z_a) + \Pi\frac{\partial}{\partial t}\left(\Psi + \ln\gamma\right) - \mathcal{H}\right\} , \tag{7}$$

where $H = \int dx\mathcal{H} = \frac{2\dot{\tau}}{\kappa}\int dx\sqrt{\gamma}$, which is the circumference functional of the circle when $\dot{\tau}$ is constant. The dynamics is that of a time-dependent system, with the time-dependence corresponding to the time-varying circumference of the circle of constant mean extrinsic curvature. As with general relativity on spatially compact manifolds in $(2+1)$ dimensions [19], the spatial metric can be chosen so that $\gamma = \gamma(t)$, i.e. $\gamma' = 0$, since any metric on a circle is globally conformal to a flat metric. Choosing a time parametrization so that $\dot{\tau}\sqrt{\gamma}$ is constant, the Hamiltonian will be time-independent.

We find after some manipulation that the canonical form of the field equations (3,4) is

$$2\Pi' - \Pi\Psi' = 0; \tag{8}$$

$$\Psi'' - \frac{1}{4}(\Psi')^2 - (\kappa\Pi)^2 + \gamma\left(\tau^2 - \Lambda/2\right) + \kappa\sum_a\sqrt{\gamma}\, m\, \delta(x - z_a(x^0)) = 0 \tag{9}$$

$$N_0'' - \dot{\tau}\gamma - N_0\left\{(\tau^2 - \Lambda/2)\gamma + \kappa/2\sum_a\sqrt{\gamma}\, m\, \delta(x - z_a(x^0))\right\} = 0 \tag{10}$$

$$N_1' - \dot{\gamma}/2 + \gamma\tau N_0 = 0 \tag{11}$$

$$\dot{\Pi} + \partial_1(-\frac{1}{\gamma}N_1\Pi + \frac{1}{2\kappa\sqrt{\gamma}}N_0\Psi' + \frac{1}{\kappa\sqrt{\gamma}}N_0') = 0 \tag{12}$$

$$\dot{\Psi} + 2N_0\left(\kappa\frac{\Pi}{\sqrt{\gamma}} + \tau\right) - N_1(\frac{1}{\gamma}\Psi') = 0 \tag{13}$$

where equilibrium solutions are characterized by $\dot{z_a} = 0 = p_a$, corresponding to a situation in which all particles are motionless at various points around the circle. A calcuation of the geodesic equations in (3) show that this implies

$$\left.\frac{\partial N_0}{\partial x}\right|_{x=z_a} = 0 \qquad\qquad N_1(z_a) = 0 \ . \tag{14}$$

Remarkably, this set of equations admits an exact solution for N bodies of equal mass m at equal time-varying proper separations. We set $c^2 \equiv \gamma\left(\tau^2 - \Lambda/2\right) = c_+^2 > 0$, and $r = |z_a - z_{a+1}| = \frac{2L}{N}$, where $a = 1, 2, \ldots, N$. The system is cyclically symmetric and so we can choose the origin to be halfway between any two particles in the N =even case or on a particle in the N =odd case. In seeking solutions to eqs. (8–13) we must ensure that the solutions for Ψ and N_0 are appropriately matched at the locations z_a of each particle and the identification point $|x| = L$; they must be continuous there but not differentiable at the locations of the particles; for example $\lim_{\epsilon\to 0}\left[\Psi'(z_a+\epsilon) - \Psi'(z_a-\epsilon)\right] = \kappa\sqrt{\gamma}m$.

With this in mind, eq. (8) is straightforwardly found to yield $\Pi = \Pi_0(t)\exp\Psi/2$, where $\Pi_0(t)$ is an arbitrary function. Upon substitution of this result into eq. (9), and employing the conditions (14), the next three equations are solved by straightforward integration and matching across the particles, furnishing a solution for Ψ, N_0, and N_1 in terms of $\Pi_0(t)$ and the metric function $\gamma(t)$. Imposing (14) and the periodicity condition $N_1(L) = N_1(-L)$ then yields a differential equation for γ whose solution is

$$\sqrt{\gamma} = \frac{N}{L\sqrt{\tau^2 - \Lambda/2}}\text{arctanh}\left(\frac{\xi\sqrt{\frac{\kappa^2 M^2}{16N^2} + (\xi^2-1)(\tau^2-\Lambda/2)} - \frac{\kappa M}{4N}}{\frac{\kappa^2 M^2}{16N^2} + \xi^2(\tau^2 - \Lambda/2)}\sqrt{\tau^2 - \Lambda/2}\right) \tag{15}$$

where $M = mN$ is the total mass of the system. The function $\Pi_0(t)$ is then found from integration of (13). Satisfaction of equation (12) provides a non-trivial consistency check on the full result, which is

$$\Psi = -2\ln\left(\beta - \frac{\sinh(\frac{c_+ L}{N})}{\sinh(c_+ L)}\sum_{a=1}^{n}\cosh(c_+(|x - z_a| - L))\right) - 2\ln(\frac{\kappa\Pi_0(t)}{c_+\sqrt{\beta^2 - 1}}) \tag{16}$$

$$N_0 = \frac{\dot{\tau}\gamma}{c_+^2\beta}\left(\frac{\sinh(\frac{c_+ L}{N})}{\sinh(c_+ L)}\sum_{a=1}^{n}\cosh(c_+(|x - z_a| - L))) - \beta\right) \tag{17}$$

$$N_1 = \gamma\frac{\dot{c_+}}{c_+}x - \frac{\gamma^2\tau\dot{\tau}}{c_+^3\beta}\frac{\sinh(\frac{c_+ L}{N})}{\sinh(c_+ L)}\sum_{a=1}^{n}\epsilon(x - z_a)\left[\sinh(c_+(|x - z_a| - L)) + \sinh(c_+ L)\right] \tag{18}$$

for the dilaton, lapse and shift functions, where the step function $\epsilon(x) = \frac{|x|}{x}$, and $\epsilon(0) = 0$. Here

$$\beta = \frac{4N}{\kappa M}\sqrt{\frac{\kappa^2 M^2}{16N^2} + (\xi^2 - 1)(\tau^2 - \Lambda/2)} \tag{19}$$

$$\Pi_0 = \sqrt{\gamma}\zeta^2\left(\tau \pm \sqrt{\frac{\kappa^2 M^2}{16N^2(\xi^2 - 1)} + (\tau^2 - \Lambda/2)}\right) \tag{20}$$

with ζ and ξ integration constants. The conjugate dilaton momentum is easily seen to be

$$\kappa\Pi = \frac{\pm c_+\sqrt{\beta^2 - 1}}{\frac{\sinh(\frac{c_+ L}{N})}{\sinh(c_+ L)}\sum_{a=1}^{n}\cosh(c_+(|x - z_a| - L)) - \beta} \tag{21}$$

We can more succinctly write the above using the relation $\sum_{a=1}^{n}\cosh(c_+(|x - z_a| - L)) = \cosh(c_+ f(x))\frac{\sinh(c_+ L)}{\sinh(\frac{c_+ L}{N})}$ where $f(x)$ is the saw-tooth function that peaks with a value of L/N (i.e. $f(z_a) = L/N$) at the particle locations and has zero magnitude half-way between the particles (i.e. $f(\frac{z_a - z_{a+1}}{2}) = 0$). This relation holds because each function is periodic, and so the origin can be shifted to be between any pair of adjacent particles. A simple shift of the origin in the cosh function and a subsequent manipulation of the sum yields the equivalence.

Another class of solutions exists for $c^2 = c_-^2 < 0$. These are obtained upon replacing $c_+ \to ic_-$ in the above results. The solution for the metric function γ now becomes

$$\sqrt{\gamma} = \frac{N}{L\sqrt{\Lambda/2 - \tau^2}}\left[\arctan\left(\frac{\xi\sqrt{\frac{\kappa^2 M^2}{16N^2} + (1 - \xi^2)(\Lambda/2 - \tau^2)} - \frac{\kappa M}{4N}}{\frac{\kappa^2 M^2}{16N^2} - \xi^2(\Lambda/2 - \tau^2)}\sqrt{\Lambda/2 - \tau^2}\right) + k\pi\right] \tag{22}$$

where k is an integer and the integration constant $|\xi| < \sqrt{1 + \frac{(\kappa M)^2}{8\Lambda N^2}}$ as a consequence of the periodicity of $N_1(x,t)$. There are no solutions for $c^2 = 0$.

The solution (15–18,21) (and its $c^2 < 0$ counterpart) is a new dynamical equilibrium solution to the N-body problem in relativistic gravity. It corresponds to an expanding/contracting spacetime of a circle with N bodies at equal time-varying proper separations from one another around the circle. For $N = 1$ the results reduce to the single particle case obtained previously [17]; indeed, the N-body solution for the spatial metric is equivalent to that of the single-particle solution with a rescaling of the parameters $L \to L/N$ and $M \to M/N$. Note that this rescaling equivalence does not hold for the remaining metric and dilaton functions.

Analysis of the behaviour of the spatial metric and Hamiltonian thus proceeds along the same lines as ref. [17]. For fixed particle mass m the spatial metric and Hamiltonian behave analogously to the single-particle case times a factor of N. For $\Lambda < 0$ the proper circumference of the circle expands from zero to some maximal size and then recontracts, whereas for $\Lambda = 0$ it undergoes a perturally decelerating expansion due to the presence of the point masses. For $\Lambda > 0$ the cosmological expansion opposes decelerating effects due to the point masses and a variety of behaviours can ensue, including expansions from zero size that asymptote to some finite value, evolutions to maximal size that begin/end at finite circumference and perpetually oscillating universes.

For fixed total mass M the evolution of the spatial metric becomes independent of M as the number of bodies N becomes large, and the full solution more closely approaches its vacuum state on a circle with $L \to L/N$.

Generalization of our approach to other dilatonic theories of gravity is straightforward. An arbitrary $(1+1)$-dimensional dilatonic theory of gravity minimally coupled to N bodies can, after conformal transformations and a rescaling of the dilaton, be written in the form (1) [4,16] with the replacement $m \to m(\psi)$. The field equations remain the same, except that (12) will contain terms proportional to $N_0\frac{\partial V}{\partial \psi}$ and $N_0\frac{\partial m}{\partial \psi}$, and $\left.\frac{\partial(mN_0)}{\partial x}\right|_{x=z_a} = 0$ will modify (14). While solving the field equations between the particles is straightforward [20], implementation of the matching conditions is not, and so the existence of an equilibrium solution in the general case remains an interesting open question.

The solution we have obtained depends quite sensitively on the matching conditions, and these in turn depend upon the equality of the point masses and their proper separations. Perturbations from this equlibrium case would be interesting to study, as they would form model inhomogeneous self-gravitating cosmological systems. For sufficiently large positive Λ, we expect that cosmological expansion will overcome the tendency to gravitationally clump, with the bodies maintaining some form of thermal equilibrium diluted by the overall expansion. For negative Λ, there should be an instability to formation of a 2d black hole [1,2].

This work was supported by the Natural Sciences and Engineering Research Council of Canada.

---

[1] R.B. Mann, Found. Phys. Lett. **4** (1991) 425; R.B. Mann, Gen. Rel. Grav. **24** (1992) 433; J.D. Brown, M. Henneaux, C. Teitelboim Phys.Rev.**D33** (1986) 319;

[2] J.D. Christensen and R.B. Mann Class.Quant.Grav. **9** (1992) 1769.

[3] T. Klösch and T. Strobl Class.Quant.Grav.**13** (1996) 965; Erratum-ibid.**14** (1997) 825.

[4] D. Louis-Martinez, J. Gegenberg and G. Kunstatter Phys.Lett. **B321** (1994) 193.

[5] G. Mandal, A.M. Sengupta and S.R. Wadia Mod.Phys.Lett. **A6** (1991) 1685.

[6] C.G. Callan, S.B. Giddings, J.A. Harvey, and A. Strominger Phys.Rev. **D45** (1992) 1005.

[7] T. Ohta and R.B. Mann, Class. Quant. Grav. **13** (1996) 2585.

[8] R.B. Mann and P. Chak, Phys.Rev.**E65** (2002) 026128.

[9] See B.N. Miller and P. Youngkins, Phys. Rev. Lett. **81** 4794 (1998); K.R. Yawn and B.N. Miller, Phys. Rev. Lett. **79** 3561 (1997) and references therein.

[10] H. Koyama and T. Kinoshi, Phys. Lett. **A** (in press; astro-ph/0008208)

[11] R.B. Mann and T. Ohta, Phys. Rev. **D55** (1997) 4723; Class. Quant. Grav. **14** (1997) 1259.

[12] R.B. Mann, D. Robbins and T. Ohta, Phys. Rev. Lett. **82** (1999) 3738.

[13] R.B. Mann, D. Robbins and T. Ohta, Phys. Rev. **D60** (1999) 104048.

[14] R.B. Mann, D. Robbins, T. Ohta and M. Trott, Nucl. Phys. **B590** 367.

[15] G. Rybicki, Astrophys. Space. Sci **14** (1971) 56.

[16] T. Banks and M. O' Loughlin, Nucl. Phys. **B362** (1991) 649; R.B. Mann, Phys. Rev.**D47** (1993) 4438.

[17] R.B. Mann, Class.Quant.Grav.**18** (2001) 3427.

[18] R. Jackiw, Nucl. Phys. B **252**, 343 (1985); C. Teitelboim, Phys. Lett. B **126**, 41, (1983).

[19] V. Moncreif, J. Math. Phys. **30** (1989) 2907.

[20] H. Pelzer and T. Strobl, Class.Quant.Grav.**15** (1998) 3803